

# Contents

Chapter 1. Introduction	1
Acknowledgments	3
Chapter 2. Examples	5
Chapter 3. Proof of Theorems 1.6 and 1.7	7
Chapter 4. Initial assumptions and reductions	9
Chapter 5. Culler-Shalen theory	11
Chapter 6. Bending characters of triangle group amalgams	15
6.1. Curves of characters of free products of cyclic groups	15
6.2. Amalgamated products	16
6.3. HNN extensions	21
Chapter 7. The proof of Theorem 1.2 when $F$ is a semi-fibre	25
7.1. Proof of Proposition 7.1 when $M(\alpha)$ is very small	25
7.2. Proof of Proposition 7.1 when $M(\alpha)$ is not very small	29
Chapter 8. The proof of Theorem 1.2 when $F$ is a fibre	45
Chapter 9. Further assumptions, reductions, and background material	59
Chapter 10. The proof of Theorem 1.2 when $F$ is non-separating but not a fibre	61
10.1. Proof of Proposition 10.1 when $M(\alpha)$ is not very small	62
10.2. Refinements to the very small case when $t_1^+ = 0$	65
Chapter 11. Algebraic and embedded $n$ -gons in $X^\epsilon$	69
Chapter 12. The proof of Theorem 1.2 when $F$ separates but is not a semi-fibre and $t_1^+ + t_1^- > 0$	75
Chapter 13. Background for the proof of Theorem 1.2 when $F$ separates and $t_1^+ = t_1^- = 0$	81
13.1. The case that $M(\beta)$ is Seifert	81
13.2. Edge weights	82
13.3. The trace of an essential annulus in a face of $\Gamma_F$	84
Chapter 14. Recognizing the figure eight knot exterior	89
14.1. Exceptional Dehn fillings of the minimally twisted chain link	89
14.2. Essential annuli in $\widehat{X^\epsilon}$ and the figure eight knot exterior	90

14.3.	The combinatorics of $\overline{\Gamma}_F$ and the figure eight knot exterior	96
Chapter 15.	Completion of the proof of Theorem 1.2 when $\Delta(\alpha, \beta) \geq 7$	103
Chapter 16.	Completion of the proof of Theorem 1.2 when $X^-$ is not a twisted $I$ -bundle	105
Chapter 17.	Completion of the proof of Theorem 1.2 when $\Delta(\alpha, \beta) = 6$ and $d = 1$	107
Chapter 18.	The case that $F$ separates but not a semi-fibre, $t_1^+ = t_1^- = 0$ , $d \neq 1$ , and $M(\alpha)$ is very small	109
Chapter 19.	The case that $F$ separates but is not a semi-fibre, $t_1^+ = t_1^- = 0$ , $d > 1$ , and $M(\alpha)$ is not very small	113
19.1.	Involutions on $M$ and its fillings	113
19.2.	Branch sets of the involutions on small Seifert manifolds	115
19.3.	The proof of Proposition 19.1	116
Chapter 20.	Proof of Theorem 1.8	119
	Bibliography	121