

CONTENTS

CHAPTER I

DEFINITIONS AND SIMPLE EXAMPLES OF GROUPS

SECT.	PAGE
1-3. Operations defined by transformation schemes; inverse operations; powers of operations; permutable operations; similar operations	1
4-7. Transformation group defined; continuous group; infinite group; discontinuous group; example of a mixed group	2
8. Identical transformation defined	3
9, 10. Examples of discontinuous groups	4
11. Examples of infinite continuous groups	4
12. Definition of finite continuous group	5
13, 14. Infinitesimal transformations; infinitesimal operators; connexion between the finite and the infinitesimal transformations of a group; effective parameters; example	6
15. Independent infinitesimal transformations; independent linear operators; unconnected operators	7
16. The alternant (X_1, X_2) defined; the alternants of the operators of a group dependent on those operators	8
17. Verification of the relation between the infinitesimal and finite transformations of a group	9
18. Simple isomorphism; example	9
19-21. The parameter groups; notation for summation	11
22. Transformation of a group; type of a group	15
23, 24. Conjugate operations; Abelian operations; operations admitted by a group; sub-group; conjugate sub-group; Abelian group; special linear homogeneous group; sub-groups of the projective group of the straight line; the translation group is the type of a group with one parameter	16
25, 26. Order of a group; projective group of the plane; its sub-groups; another type of group; projective group of space; some of its sub-groups; similar groups	18
27. Euler's transformation formulae	20
28. A non-projective group; number of types increases with the number of variables	22

CHAPTER II

ELEMENTARY ILLUSTRATIONS OF THE PRINCIPLE OF
EXTENDED POINT TRANSFORMATIONS

SECT.	PAGE
29. Differential equations admitting known groups . . .	23
30. The extended infinitesimal point transformation . . .	23
31, 32. Differential invariants; example; some particular classes of differential equations admit infinitesimal point transformations; examples	25
33. Relation between the equations $\left(\frac{\partial u}{\partial x}\right)^2 + \left(\frac{\partial u}{\partial y}\right)^2 + \left(\frac{\partial u}{\partial z}\right)^2 = 0$ and $dx^2 + dy^2 + dz^2 = 0$; minimum curves; Mongian equations	28
34. Direct verification of this relation	29
35, 36. The infinitesimal operators admitted by lines of zero length; the conformal transformation group	31

CHAPTER III

THE GENERATION OF A GROUP FROM ITS INFINITESIMAL
TRANSFORMATIONS

37, 38. The parameters which define the identical transforma- tion; example illustrating method of finding these parameters; the symbol ϵ_{hi}	34
39. A group not containing the identical transformation . . .	36
40. Method of obtaining the infinitesimal operators	36
41-43. First fundamental theorem; the order of a group equals the number of its operators; illustrative examples	38
44. The operators of the first parameter group; preliminary formulae and proof of the theorem that <i>every finite operation of a group may be generated by indefinite repetition of an infinitesimal one</i> ; transitive group; simply transitive group; canonical equations of a group; simple relation between an operation and its inverse when the group is in canonical form	41
45, 46. On finding the finite equations of a group when the infinitesimal operators are given; example; the finite equations of the group not often required	46

CHAPTER IV

THE CONDITIONS THAT A GIVEN SET OF LINEAR OPERATORS
MAY GENERATE A GROUP

SECT.	PAGE
47. The fundamental set of operators of a group not unique; the structure constants; the second fundamental theorem stated and proved; the converse stated; illustrative example	51
48, 49. Formal laws of combination of linear operators; examples	53
50, 51. Proof of converse of the second fundamental theorem	57
52. Form in which any operation of the group can be expressed	59
53, 54. Examples	60
55. Reciprocal groups; example	62
56, 57. Examples	62
58. The structure constants of a group the same as those of its first parameter group	65
59. The converse of the first fundamental theorem	65

CHAPTER V

THE STRUCTURE CONSTANTS OF A GROUP

60. Jacobi's identity; relation between the structure constants of a group; the third fundamental theorem; statement of its converse	67
61. The structure constants vary with the choice of the fundamental sets of operators; groups of the same structure	68
62. The normal structure constants	70
63. The group $X_i = \sum_{j=k=n} c_{jik} x_j \frac{\partial}{\partial x_k}$; example on group construction	73
64. Proof of converse of third fundamental theorem; example on group construction	74
65, 66. Solution of a system of differential equations	76
67. The three fundamental theorems	80

CHAPTER VI

COMPLETE SYSTEMS OF DIFFERENTIAL EQUATIONS

SECT.	PAGE
68. The terms unconnected and independent distinguished ; complete system of operators ; functions admitting infinitesimal transformations	81
69. The invariants of a complete system ; complete system in normal form	83
70, 71. The form to which a complete system in normal form can be reduced	84
72. The number of invariants of a complete system ; how obtained	87
73. Integration operations of given order defined ; how the order of the operations necessary for the solution of a given equation is lowered, when the given equation is a member of a given complete system ; an operator which annihilates the invariants of a complete system belongs to that system	87

CHAPTER VII

DIFFERENTIAL EQUATIONS ADMITTING KNOWN TRANSFORMATION GROUPS

74. Object of the chapter ; formula for the transformation of an operator to new variables	90
75. Condition that a sub-group may be self-conjugate	91
76, 77. Condition that a complete system of equations may admit an infinitesimal transformation ; second form of the condition	93
78. Trivial transformations ; distinct transformations ; condition that a system admitting $x' = x + t\xi$ may also admit $x' = x + t\rho\xi$	95
79. Reduced operators ; if a system admits any operators it admits a complete system of operators	96
80. Properties of such a complete system ; it may give some integrals of the given complete system of equations	97
81. Form of the admitted operators necessary for further advance towards solution of system	98

SECT.	PAGE
82. Restatement of problem at this stage; origin of Lie's group-theory	99
83. Simplification and further restatement	100
84. Maximum sub-group of group admitted; new integrals .	101
85. If this sub-group is not self-conjugate further integrals may be obtained without integration operations . . .	101
86. Completion of the proposed problem	104
87. All the integrals can be obtained by quadratures when the group is of a certain form	105
88-92. General remarks on the foregoing theory; application to examples; points of special position; theorem about these points stated; further examples	106

CHAPTER VIII

INVARIANT THEORY OF GROUPS

93-95. Second definition of transitivity; invariants of intransitive group; geometrical interpretation; cogredient transformation; groups extended into point-pair groups	113
96, 97. The invariant theory of algebra in relation to group-theory	116
98-100. The functional form which in the more general invariant theory takes the place of the quantic in the invariant theory of algebra; the invariant theory of this form; how the form may be obtained; example	119
101. Proof of theorem as to points of special position	124
102-104. Invariant equations with respect to a group; contracted operators of the group with respect to these equations; proof of formula $(\overline{X}_k \overline{f}) \equiv \overline{X}_k . \overline{f}$; contracted operators generate a group; the order of the special points defined by the invariant equations is equal to the number of unconnected contracted operators	127
105, 106. Equations which admit the infinitesimal transformations admit all the transformations of the group; method of obtaining such equations; examples	130

CHAPTER IX

PRIMITIVE AND STATIONARY GROUPS

107. Geometrical interpretation of the invariants of an intransitive group; the contracted operators, with respect to any invariant manifold of the group	135
---	-----

SECT.	PAGE
108-110. Primitive and imprimitive groups; the group which transforms a manifold of an imprimitive group into some other such manifold; groups admitted by a complete system of differential equations are imprimitive, and conversely	136
111, 112. The sub-group of a point; the operators of this sub-group; conditions that the sub-groups of two points may be coincident	139
113, 114. Stationary and non-stationary groups; analytical proof that a stationary group is imprimitive	141
115. The functions $\phi_{j\mu}$, Π_{ijk} ; structure functions; stationary functions	143
116. Simplification of stationary functions; the group Z_1, \dots, Z_r can be constructed when the structure constants and stationary functions are assigned	144
117. Reduction of the operators of a group to standard form	145
118. Integration operations necessary to find the finite equations of a stationary group	147

CHAPTER X

CONDITION THAT TWO GROUPS MAY BE SIMILAR. RECIPROCAL GROUPS

119, 120. Necessary conditions for similarity; simplification of these conditions preparatory to proving that they are also sufficient	148
121. A correspondence between a q -fold in x space and a q -fold in y space; initial points; the general correspondence between the two spaces; proof that it is a point-to-point correspondence	151
122. Proof that the necessary conditions for similarity are sufficient	153
123, 124. If s is the number of unconnected stationary functions there are $(n-s)$ unconnected operators Z_1, \dots, Z_{n-s} permutable with each of the operators X_1, \dots, X_r ; Z_1, \dots, Z_{n-s} form a complete system whose structure functions are invariants of X_1, \dots, X_r	154
125. If X_1, \dots, X_r is transitive, Z_1, \dots, Z_{n-s} are the operators of a group; if simply transitive, Z_1, \dots, Z_{n-s} is also simply transitive, and has the same structure constants as X_1, \dots, X_r	157

CHAPTER XI

ISOMORPHISM

SECT.	PAGE
126, 127. The operators of the parameter groups in canonical form expanded for a few terms in powers of the variables; direct proof that the structure constants of a group and its first parameter group are the same; the canonical form of a group not fixed till the fundamental set of operators is chosen	159
128. Two groups simply isomorphic when they have the same parameter group	162
129, 130. When one group is multiply isomorphic with another; a self-conjugate sub-group within the first corresponds to the identical transformation in the second; condition for isomorphic relation between two groups; simple groups	162
131, 132. When the structure constants of a group are given, the structure constants of every group with which the first is multiply isomorphic can be found; the isomorphic relation which may exist between the r independent operators of a group, and the r non-independent operators of a group whose order is less than r	165
133. Examples of groups isomorphically related; proof that two transitive groups in the same number of variables are similar, if they are simply isomorphic in such a way that the sub-group of some point of general position in the one corresponds to the sub-group of some point of general position in the other	167

CHAPTER XII

ON THE CONSTRUCTION OF GROUPS WHOSE STRUCTURE CONSTANTS AND STATIONARY FUNCTIONS ARE KNOWN

134. Object of the chapter	169
135. General relation between the structure functions of any complete system of operators; simplification of the problem to be discussed	170
136. The system of simultaneous differential equations on whose solution the problem depends	171
137. Proof that this system is a consistent one; general method of solution	173
138. Extension so as to apply to the case of intransitive groups	174

CHAPTER XIII

CONJUGATE SUB-GROUPS: THE CONSTRUCTION OF GROUPS
FROM THEIR STRUCTURE CONSTANTS

SECT.	PAGE
139. A new set of fundamental operators Y_1, \dots, Y_r is chosen instead of the original set X_1, \dots, X_r	176
140, 141. Definition of the functions H_{ijk} ; proof that they are the structure constants of Y_1, \dots, Y_r ; the functions Π_{ijk} ; identity connecting these functions	177
142, 143. Definition of the operators Π_1, \dots, Π_r ; they form a group with which X_1, \dots, X_r is isomorphic; the equation system $H_{q+i, q+j, k} = 0$ admits these operators	179
144, 145. The equation system $H_{q+i, q+j, k} = 0$ defines sub-groups of order $r - q$; method of finding all such sub-groups; the group within which a given sub-group is invariant; the index of a sub-group	181
146. Method of finding all sub-groups conjugate to a given sub-group	183
147. Method of finding all the different types of sub-groups	186
148. Application of the preceding discussion to enable us to determine the stationary functions of a group whose structure constants are given	187
149-151. Illustrative examples; a particular case of the general theory	189

CHAPTER XIV

ON PFAFF'S EQUATION AND THE INTEGRALS OF PARTIAL
DIFFERENTIAL EQUATIONS

152. Element of space; united elements; Pfaff's equation and its solution; Pfaffian system of any order; generating equations	194
153, 154. Alternant of two functions; functions in involution; equations in involution; homogeneous function system; necessary and sufficient conditions that n equations should form a Pfaffian system	196
155. Geometrical interpretation of solution of Pfaff's equation	201
156, 157. Lie's definition of an integral; the problem involved in the solution of a partial differential equation of the first order	202

SECT.	PAGE
158. Proof that $(\bar{u}, \bar{v}) \equiv \overline{(u, v)}$	205
159. Proof that P is not connected with $\bar{u}_1, \dots, \bar{u}_m$	206
160-163. On finding the complete integral of a given equation; illustrative examples on the foregoing theory	208

CHAPTER XV

COMPLETE SYSTEMS OF HOMOGENEOUS FUNCTIONS

164. Necessary and sufficient conditions that a given system of functions may be a homogeneous one	213
165. General definition of a complete homogeneous function system; structure functions of the system; if all the functions are of zero degree the system is in involution	214
166. If f is annihilated by $\bar{u}_1, \dots, \bar{u}_m$, where u_1, \dots, u_m form a homogeneous function system, Pf is also annihilated	215
167. Proof of the identity $(u, (v, w)) + (w, (u, v)) + (v, (w, u)) \equiv 0$; the polar system	216
168. The functions common to a system and its polar are homogeneous and in involution thus forming an Abelian sub-system; satisfied system	217
169, 170. Any complete homogeneous system is a sub-system within a satisfied system; complete systems of the same structure; contracted operator of \bar{u}_i	218
171-174. The normal forms of complete homogeneous systems; systems of the same structure	220
175, 176. Every complete system of homogeneous functions is a sub-system within a system of order $2n$; two systems of the same structure are sub-systems of the same structure within two systems of the same structure and of order $2n$	223

CHAPTER XVI

CONTACT TRANSFORMATIONS

177. Necessary and sufficient conditions that $x'_i = X_i, p'_i = P_i$ should lead to $\sum_{i=1}^n p'_i dx'_i = \sum_{i=1}^n p_i dx_i$	226
178, 179. $X_1, \dots, X_n, P_1, \dots, P_n$ are unconnected; contact transformation defined; geometrical interpretation; the transformation given when X_1, \dots, X_n are given; example of contact transformation	228

SECT.	PAGE
180, 181. By a contact transformation a Pfaffian system of equations is transformed into a Pfaffian system; examples on the application of contact transformations to differential equations	231
182, 183. Any two complete homogeneous systems of functions of the same structure, and in the same number of variables, can be transformed into one another by a homogeneous contact transformation; extension to the case of non-complete systems of functions	234
184, 185. Non-homogeneous form of Pfaff's equation; the corresponding Pfaffian systems and contact transformations	238
186, 187. Example on the reduction of a function group to a simple form by a contact transformation; Ampère's equation reducible to the form $s = 0$ if it admits two systems of intermediary integrals	241

CHAPTER XVII

THE GEOMETRY OF CONTACT TRANSFORMATIONS

188. The generating equations of a contact transformation	245
189, 190. Limitation on the form which generating equation can assume; interpretation	246
191. Contact transformation with a single generating equation	247
192. Special elements, and the special envelope	249
193, 194. The three classes of element manifolds	250
195. Reciprocation	252
196, 197. Contact transformation with two generating equations	253
198. Linear complexes	255
199, 200. Bilinear equations as generating equations, simplification	257
201, 202. The generating equations $x' + iy' + xz' + z = 0, \quad x(x' - iy') - y - z' = 0;$ to points in space x, y, z correspond minimum lines in x', y', z' ; to points in space x', y', z' correspond lines of a linear complex in x, y, z	259
203-205. To lines in space x, y, z , spheres in x', y', z' ; to spheres in x', y', z' , a positive and a negative correspondent in x, y, z ; contact of spheres and intersection of lines; example	262
206-210. To a quadric in x, y, z a cyclide in x', y', z' ; to lines of inflexion, lines of curvature; further examples	265

SECT.	PAGE
211-217. The generating equations $axx' + byy' + czz' + d = 0, \quad xx' + yy' + zz' + 1 = 0$ transform a point in one space to a line of the tetrahedral complex in the other; a plane to a twisted cubic; a straight line to a quadric; deduction of geometrical theorem; the generators in the quadric; case of degeneration; illustrative examples on method	268
218. Point transformation	275

CHAPTER XVIII

INFINITESIMAL CONTACT TRANSFORMATIONS

219-221. Infinitesimal contact transformations; characteristic functions; condition that an equation should admit an infinitesimal contact transformation	276
222, 223. Characteristic manifolds of an equation; transformation of, by a contact transformation; geometrical interpretation of infinitesimal contact transformation	278
224, 225. Linear element; elementary integral cone; Mongian equations; correspondence between Mongian equations and partial differential equations; Mongian equations and partial differential equation related to tetrahedral complex	280
226, 227. The characteristic function of the alternant of two contact operators; transformation of operator by a given contact transformation	284
228-232. Finite contact group; extended point group; its structure; condition that two contact groups may be similar	286
233, 234. Reducible contact groups; contact groups regarded as point groups in space of higher dimensions	292

CHAPTER XIX

THE EXTENDED INFINITESIMAL CONTACT TRANSFORMATIONS: APPLICATIONS TO GEOMETRY

235, 236. The transformation of the higher derivatives of z by an infinitesimal contact transformation; explicit forms for $\frac{d^2y}{dx^2}$ and $\frac{d^3y}{dx^3}$	294
237-239. The groups transforming straight lines into straight lines, circles into circles	297

SECT.	PAGE
240, 241. Transformations of the group of § 239; correspondence between circles of the plane and lines of a linear complex in space; a projective group isomorphic with the conformal group	302
242, 243. The twice extended contact operator in three variables; transformations admitted by $s=0$; Ampère's equation	305
244, 245. The transformations which do not alter the length of arcs on a given surface; the measure of curvature unaltered by such	308
246-249. Surfaces over which a net can move; geometrical treatment of the question; analytical discussion by aid of Gaussian coordinates; the group of movements of the net	311

CHAPTER XX

DIFFERENTIAL INVARIANTS

250, 251. How to obtain the differential invariants of a given group	319
252. Differential invariants of the group $x' = x, y' = \frac{ay+b}{cy+d}$.	321
253, 254. Extended operators of projective group of the plane; invariant differential equations; absolute differential invariants; the invariants of lowest order of this group	322
255-257. The group of movements in non-Euclidean space; extended operators of; differential invariants of; geometrical considerations help in determination of . .	326

CHAPTER XXI

THE GROUPS OF THE STRAIGHT LINE, AND THE PRIMITIVE GROUPS OF THE PLANE

258. The possible types of groups in a given number of variables	331
259. Operators arranged in systems according to degree of the coefficients in the variables	332
260. The possible types of groups in a single variable	333
261, 262. Simplification of any operator of the linear homogeneous group	335

SECT.	PAGE
263, 264. The possible types of linear homogeneous groups in the plane	339
265-270. The primitive groups of the plane; operators of the first degree; the group cannot have operators of the third degree; possible form of operators of the second degree; structure constants of the group; possible types of .	342

CHAPTER XXII

THE IMPRIMITIVE GROUPS OF THE PLANE

271. Can be arranged in four classes, and thus successively found	353
272-274. The groups of the first class	354
275-279. The groups of the second class	357
280, 281. The groups of the third class	362
282. The groups of the fourth class	364
283. The systems of curves which are invariant for the different types of imprimitive groups	365
284. Enumeration of the mutually exclusive types of imprimitive groups of the plane	368

CHAPTER XXIII

THE IRREDUCIBLE CONTACT TRANSFORMATION GROUPS OF THE PLANE

285-287. Condition for the reducibility of a system of contact operators of the plane; an irreducible group of the plane is a transitive group of space; the form of the operators of the first degree	370
288-290. The irreducible groups in the first class have six independent operators; the structure of any such group . . .	373
291, 292. Every group in this class is of the same type . . .	377
293, 294. The remaining irreducible contact groups of the plane .	378

CHAPTER XXIV

THE PRIMITIVE GROUPS OF SPACE

295, 296. The curves which admit two infinitesimal projective transformations must be straight lines or conics . . .	381
297. Any sub-group of the projective group must leave unaltered either a point, a line, or a conic	383

SECT.	PAGE
298. A projective group isomorphic with the group of the origin; the cases when this projective group has no invariant	385
299, 300. The case when it has as invariant a straight line . . .	386
301-305. The cases when it has as invariant a conic	389
306. Enumeration of the types of primitive groups	396

CHAPTER XXV

SOME LINEAR GROUPS CONNECTED WITH HIGHER COMPLEX NUMBERS

307-309. Properties of simply transitive groups which involve the variables and the parameters linearly in their finite equations	398
310, 311. Determination of all the groups of this class in three variables	401
312. The theory of higher complex numbers	406
313-315. To every such system a group of the class considered will correspond, and conversely. Examples	408
INDEX	411