

# CONTENTS

<b>Chapter 1</b>			
<b>An Invitation to Neurobiology</b>	<b>1</b>		
NATURE AND NURTURE IN BRAIN FUNCTION AND BEHAVIOR	1		
1.1 Human twin studies can reveal the contributions of nature and nurture	1		
1.2 Examples of nature: Animals exhibit instinctive behaviors	3		
1.3 An example of nurture: Barn owls adjust their auditory map to match an altered visual map	4		
HOW IS THE NERVOUS SYSTEM ORGANIZED?	6		
1.4 The nervous system consists of neurons and glia	7		
1.5 Individual neurons were first visualized by Golgi staining in the late nineteenth century	8		
1.6 Twentieth-century technology confirmed the neuron doctrine	10		
1.7 In vertebrate neurons, information generally flows from dendrites to cell bodies to axons	11		
1.8 Neurons use membrane potential changes and neurotransmitter release to transmit information	13		
1.9 Neurons function in the context of specific neural circuits	15		
1.10 Specific brain regions perform specialized functions	18		
1.11 The brain uses maps to organize information	19		
1.12 The brain is a massively parallel computational device	21		
GENERAL METHODOLOGY	23		
1.13 Observations and measurements are the foundations for discovery	23		
1.14 Perturbation experiments establish causes and mechanisms	24		
SUMMARY	25		
FURTHER READING	25		
<b>Chapter 2</b>			
<b>Signaling within Neurons</b>	<b>27</b>		
CELL BIOLOGICAL AND ELECTRICAL PROPERTIES OF NEURONS	28		
2.1 Neurons follow the central dogma of molecular biology and rules of intracellular vesicle trafficking	28	2.2 While some dendritic and axonal proteins are synthesized from mRNAs locally, most are actively transported from the soma	30
		2.3 The cytoskeleton forms the basis of neuronal polarity and directs intracellular trafficking	32
		2.4 Channels and transporters move solutes passively or actively across neuronal membranes	34
		2.5 Neurons are electrically polarized at rest because of ion concentration differences across the plasma membrane and differential ion permeability	38
		2.6 Neuronal plasma membrane can be described in terms of electrical circuits	40
		2.7 Electrical circuit models can be used to analyze ion flows across glial and neuronal plasma membrane	43
		2.8 Passive electrical properties of neurons: electrical signals evolve over time and decay across distance	44
		2.9 Active electrical properties of neurons: depolarization above a threshold produces action potentials	47
		HOW DO ELECTRICAL SIGNALS PROPAGATE FROM THE NEURONAL CELL BODY TO ITS AXON TERMINALS?	49
		2.10 Action potentials are initiated by depolarization-induced inward flow of Na <sup>+</sup>	49
		2.11 Sequential, voltage-dependent changes in Na <sup>+</sup> and K <sup>+</sup> conductances account for action potentials	50
		2.12 Action potentials are all-or-none, are regenerative, and propagate unidirectionally in the axon	52
		2.13 Action potentials propagate more rapidly in axons with larger diameters and in myelinated axons	53
		2.14 Patch clamp recording enables the study of current flow across individual ion channels	57
		2.15 Cloning of genes that encode ion channels allows their structure–function relationship to be studied	59
		2.16 Crystal structures reveal the atomic bases of ion channel properties	62
		SUMMARY	65
		FURTHER READING	66

**Chapter 3**  
**Signaling across Synapses 69**

**HOW IS NEUROTRANSMITTER RELEASE CONTROLLED AT THE PRESYNAPTIC TERMINAL? 69**

- 3.1 Action potential arrival at the presynaptic terminal triggers neurotransmitter release 69
- 3.2 Neurotransmitters are released in discrete packets 70
- 3.3 Neurotransmitters are released when synaptic vesicles fuse with the presynaptic plasma membrane 72
- 3.4 Neurotransmitter release is controlled by Ca<sup>2+</sup> entry into the presynaptic terminal 74
- 3.5 SNARE and SM proteins mediate synaptic vesicle fusion 75
- 3.6 Synaptotagmin serves as a Ca<sup>2+</sup> sensor to trigger synaptic vesicle fusion 78
- 3.7 The presynaptic active zone is a highly organized structure 79
- 3.8 Neurotransmitters are efficiently cleared from the synaptic cleft by enzymatic cleavage or transport into presynaptic and glial cells 80
- 3.9 Synaptic vesicle recycling by endocytosis is essential for continual synaptic transmission 81
- 3.10 Synapses can be facilitating or depressing 83
- 3.11 The nervous system uses many neurotransmitters 85

**HOW DO NEUROTRANSMITTERS ACT ON POSTSYNAPTIC NEURONS? 87**

- 3.12 Acetylcholine opens a nonselective cation channel at the neuromuscular junction 88
- 3.13 The skeletal muscle acetylcholine receptor is a ligand-gated ion channel 90
- 3.14 Neurotransmitter receptors are ionotropic or metabotropic 91
- 3.15 AMPA and NMDA glutamate receptors are activated by glutamate under different conditions 93
- 3.16 The postsynaptic density is organized by scaffolding proteins 95
- 3.17 Ionotropic GABA and glycine receptors are Cl<sup>-</sup> channels that mediate inhibition 96
- 3.18 All metabotropic neurotransmitter receptors trigger G protein cascades 99
- 3.19 A GPCR signaling paradigm: β-adrenergic receptors activate cAMP as a second messenger 100
- 3.20 α and βγ G protein subunits trigger diverse signaling pathways that alter membrane conductance 102

- 3.21 Metabotropic receptors can act on the presynaptic terminal to modulate neurotransmitter release 104
- 3.22 GPCR signaling features multiple mechanisms of signal amplification and termination 106
- 3.23 Postsynaptic depolarization can induce new gene expression 106
- 3.24 Dendrites are sophisticated integrative devices 110
- 3.25 Synapses are strategically placed at specific locations in postsynaptic neurons 113

**SUMMARY 116**

**FURTHER READING 118**

**Chapter 4**  
**Vision 121**

**HOW DO RODS AND CONES DETECT LIGHT SIGNALS? 121**

- 4.1 Psychophysical studies revealed that human rods can detect single photons 122
- 4.2 Electrophysiological studies identified the single-photon response of rods: light hyperpolarizes vertebrate photoreceptors 123
- 4.3 Light activates rhodopsin, a prototypical G-protein-coupled receptor 124
- 4.4 Photon-induced signals are greatly amplified by a transduction cascade 125
- 4.5 Light-triggered decline of cyclic-GMP level directly leads to the closure of cation channels 126
- 4.6 Recovery enables the visual system to respond to light continually 127
- 4.7 Adaptation enables the visual system to detect contrast over a wide range of light levels 129
- 4.8 Cones are concentrated in the fovea for high-acuity vision 130
- 4.9 Cones are less sensitive but faster than rods 131
- 4.10 Photoreceptors with different spectral sensitivities are needed to sense color 132
- 4.11 Humans have three types of cones 133
- 4.12 Cloning of the cone opsin genes revealed the molecular basis of color detection 134
- 4.13 Defects in cone opsin genes cause human color blindness 135

**HOW ARE SIGNALS FROM RODS AND CONES ANALYZED IN THE RETINA? 135**

- 4.14 Retinal ganglion cells use center-surround receptive fields to analyze contrast 136

4.15	Bipolar cells are either depolarized or hyperpolarized by light based on the glutamate receptors they express	137	5.4	Gradients of ephrins and Eph receptors instruct retinotectal mapping	172
4.16	Lateral inhibition from horizontal cells constructs the center-surround receptive fields	138	5.5	A single gradient is insufficient to specify an axis	174
4.17	Diverse retinal cell types and their precise connections enable parallel information processing	140	5.6	To cross, or not to cross: that is the question	178
4.18	Direction-selectivity of RGCs arises from asymmetric inhibition by amacrine cells	142	<b>HOW DO EXPERIENCE AND NEURONAL ACTIVITY CONTRIBUTE TO WIRING?</b>		180
4.19	Color is sensed by comparing signals from cones with different spectral sensitivities	143	5.7	Monocular deprivation markedly impairs visual cortex development	180
4.20	The same retinal cells and circuits can be used for different purposes	145	5.8	Competing inputs are sufficient to produce spatial segregation at the target	182
<b>HOW IS INFORMATION PROCESSED IN THE VISUAL CORTEX?</b>		146	5.9	Ocular dominance columns in V1 and eye-specific layers in LGN develop by gradual segregation of eye-specific inputs	183
4.21	Retinal information is topographically represented in the lateral geniculate nucleus and visual cortex	146	5.10	Retinal neurons exhibit spontaneous waves of activity before the onset of vision	184
4.22	Receptive fields of LGN neurons are similar to those of RGCs	148	5.11	Retinal waves and correlated activity drive segregation of eye-specific inputs	185
4.23	Primary visual cortical neurons respond to lines and edges	149	5.12	Hebb's rule: correlated activity strengthens synapses	187
4.24	How do visual cortical neurons acquire their receptive fields?	150	5.13	A Hebbian molecule: the NMDA receptor acts as a coincidence detector	189
4.25	Cells with similar properties are vertically organized in the visual cortex	151	<b>HOW DO MOLECULAR DETERMINANTS AND NEURONAL ACTIVITY WORK TOGETHER?</b>		190
4.26	Information generally flows from layer 4 to layers 2/3 and then to layers 5/6 in the neocortex	154	5.14	Ephrins and retinal waves act in parallel to establish the precise retinocollicular map	192
4.27	Visual information is processed in parallel streams	157	5.15	Ephrins and retinal waves also work together to establish the retinotopic map in the visual cortex	193
4.28	Face recognition cells form a specialized network in the primate temporal cortex	159	5.16	Different aspects of visual system wiring rely differentially on molecular cues and neuronal activity	195
4.29	Linking perception to decision and action: microstimulation of MT neurons biased motion choice	160	<b>VISUAL SYSTEM DEVELOPMENT IN DROSOPHILA: LINKING CELL FATE TO WIRING SPECIFICITY</b>		197
SUMMARY		163	5.17	Cell-cell interactions determine photoreceptor cell fates: R7 as an example	198
FURTHER READING		164	5.18	Multiple parallel pathways participate in layer-specific targeting of R8 and R7 axons	201
<b>Chapter 5</b>			<b>SUMMARY</b>		203
<b>Wiring of the Visual System</b>		167	<b>FURTHER READING</b>		204
<b>HOW DO RETINAL GANGLION CELL AXONS FIND THEIR TARGETS?</b>		167	<b>Chapter 6</b>		
5.1	Optic nerve regeneration experiments suggested that RGC axons are predetermined for wiring	168	<b>Olfaction, Taste, Audition, and Somatosensation</b>		207
5.2	Point-to-point connections between retina and tectum arise by chemoaffinity	169	<b>HOW DO WE SENSE ODORS?</b>		207
5.3	The posterior tectum repels temporal retinal axons	171	6.1	Odorant binding leads to opening of a cyclic nucleotide-gated channel in olfactory receptor neurons	208

6.2	<i>Ca</i> <sup>2+</sup> coordinates olfactory recovery and adaptation	210	AUDITION: HOW DO WE HEAR AND LOCALIZE SOUNDS?	238	
6.3	Odorants are represented by combinatorial activation of olfactory receptor neurons	210	6.22	Sounds are converted to electrical signals by mechanically gated ion channels in the stereocilia of hair cells	239
6.4	Odorant receptors are encoded by many hundreds of genes in mammals	210	6.23	Sound frequencies are represented as a tonotopic map in the cochlea	240
6.5	Polymorphisms in odorant receptor genes contribute to individual differences in odor perception	213	6.24	Motor properties of outer hair cells amplify auditory signals and sharpen frequency tuning	243
6.6	Each olfactory receptor neuron (ORN) expresses a single odorant receptor	214	6.25	Auditory signals are processed by multiple brainstem nuclei before reaching the cortex	245
6.7	ORNs expressing a given odorant receptor are broadly distributed in the nose	214	6.26	In the owl, sound location is determined by comparing the timing and levels of sounds reaching two ears	246
6.8	ORNs expressing the same odorant receptor project their axons to the same glomerulus	215	6.27	Mechanisms of sound location in mammals differ from those in the owl	249
6.9	Olfactory bulb circuits transform odor representation through lateral inhibition	217	6.28	The auditory cortex analyzes complex and biologically important sounds	250
6.10	Olfactory inputs are differentially organized in distinct cortical areas	218	SOMATOSENSATION: HOW DO WE SENSE BODY MOVEMENT, TOUCH, TEMPERATURE, AND PAIN?	255	
HOW DO WORMS AND FLIES SENSE ODORS?		222	6.29	Many types of sensory neurons are used to encode diverse somatosensory stimuli	257
6.11	<i>C. elegans</i> encodes olfactory behavioral choices at the sensory neuron level	223	6.30	Merkel cells and some touch sensory neurons employ Piezo2 as a mechanotransduction channel	259
6.12	<i>C. elegans</i> sensory neurons are activated by odorant withdrawal and engage ON- and OFF-pathways	224	6.31	TRP channels are major contributors to temperature, chemical, and pain sensation	262
6.13	The olfactory systems in insects and mammals share many similarities	225	6.32	Sensation can be a product of central integration: the distinction of itch and pain as an example	264
6.14	The antennal lobe transforms ORN input for more efficient representation by projection neurons	226	6.33	Touch and pain signals are transmitted by parallel pathways to the brain	266
6.15	Odors with innate behavioral significance use dedicated olfactory processing channels	230	6.34	Pain is subjected to peripheral and central modulation	268
6.16	Odor representation in higher centers is stereotyped or stochastic depending on whether the center directs innate or learned behavior	231	6.35	Linking neuronal activity with touch perception: from sensory fiber to cortex	269
TASTE: TO EAT, OR NOT TO EAT?		232	SUMMARY	272	
6.17	Mammals have five classic taste modalities: sweet, bitter, umami, salty, and sour	233	FURTHER READING	273	
6.18	Sweet and umami are sensed by heterodimers of the T1R family of G-protein-coupled receptors	233	<b>Chapter 7</b>		
6.19	Bitter is sensed by a family of ~30 T2R G-protein-coupled receptors	234	<b>Wiring of the Nervous System</b>	<b>277</b>	
6.20	Sour and salty tastes involve specific ion channels	236	HOW DOES WIRING SPECIFICITY ARISE IN THE DEVELOPING NERVOUS SYSTEM?	278	
6.21	Activation of specific taste receptor cells confers specific taste perceptions	236	7.1	The nervous system is highly patterned as a consequence of early developmental events	278
			7.2	Orderly neurogenesis and migration produce many neuronal types that occupy specific positions	280

7.3	Cell fates are diversified by asymmetric cell division and cell–cell interactions	281	HOW DO ~20,000 GENES SPECIFY 10 <sup>14</sup> CONNECTIONS?	316	
7.4	Transcriptional regulation of guidance molecules links cell fate to wiring decision	283	7.24	Some genes can produce many protein variants	316
7.5	Crossing the midline: Combinatorial actions of guidance receptors specify axon trajectory choice	286	7.25	Protein gradients can specify different connections	318
7.6	Crossing the midline: Axons switch responses to guidance cues at intermediate targets	288	7.26	The same molecules can serve multiple functions	318
7.7	The cell polarity pathway participates in determining whether a neuronal process becomes an axon or a dendrite	290	7.27	The same molecules can be used at multiple times and places	318
7.8	Local secretory machinery is essential for dendrite morphogenesis and microtubule organization	292	7.28	Combinatorial use of wiring molecules can reduce the number of wiring molecules needed	319
7.9	Homophilic repulsion enables self-avoidance of axonal and dendritic branches	293	7.29	Dividing wiring decisions into multiple steps can conserve molecules and increase fidelity	319
7.10	Subcellular site selection of synaptogenesis uses both attractive and repulsive mechanisms	295	7.30	Many connections do not need to be specified at the level of individual synapses or neurons	320
7.11	Bidirectional trans-synaptic communication directs the assembly of synapses	297	7.31	Wiring can be instructed by neuronal activity and experience	320
7.12	Astrocytes stimulate synapse formation and maturation	299	SUMMARY	321	
7.13	Activity and competition refine neuromuscular connectivity	300	FURTHER READING	322	
7.14	Developmental axon pruning refines wiring specificity	301	<b>Chapter 8</b>		
7.15	Neurotrophins from target cells support the survival of sensory, motor, and sympathetic neurons	302	<b>Motor and Regulatory Systems</b>	<b>325</b>	
ASSEMBLY OF OLFACTORY CIRCUITS: HOW DO NEURAL MAPS FORM?		305	HOW IS MOVEMENT CONTROLLED?	326	
7.16	Neural maps can be continuous, discrete, or a combination of the two	305	8.1	Muscle contraction is mediated by sliding of actin and myosin filaments and is regulated by intracellular Ca <sup>2+</sup>	326
7.17	In mice, odorant receptors instruct ORN axon targeting by regulating expression of guidance molecules	307	8.2	Motor units within a motor pool are recruited sequentially from small to large	329
7.18	ORN axons sort themselves by repulsive interactions before reaching their target	309	8.3	Motor neurons receive diverse and complex input	330
7.19	Activity-dependent regulation of adhesion and repulsion refines glomerular targeting	310	8.4	Central pattern generators coordinate rhythmic contraction of muscles during locomotion	332
7.20	<i>Drosophila</i> projection neurons' lineage and birth order specify the glomeruli that their dendrites target	312	8.5	Intrinsic properties of neurons and their connection patterns produce rhythmic output in a model central pattern generator	334
7.21	Graded determinants and discrete molecular labels control the targeting of projection neuron dendrites	313	8.6	The spinal cord uses multiple central pattern generators to control locomotion	336
7.22	Sequential interactions among ORN axons limit their target choice	314	8.7	The brainstem contains specific motor control nuclei	338
7.23	Homophilic matching molecules instruct connection specificity between synaptic partners	315	8.8	The cerebellum is required for fine control of movement	340
			8.9	The basal ganglia participate in initiation and selection of motor programs	343
			8.10	Voluntary movement is controlled by the population activity of motor cortical neurons in a dynamical system	346

8.11	Population activity of motor cortical neurons can be used to control neural prosthetic devices	349	9.2	<i>Fruitless</i> ( <i>Fru</i> ) is essential for many aspects of sexual behavior	379	
<b>HOW DOES THE BRAIN REGULATE THE FUNCTIONS OF INTERNAL ORGANS?</b>			351	9.3	A sex-determination hierarchy specifies sex-specific splicing of <i>Fru</i> that produces male-specific <i>Fru<sup>M</sup></i>	379
8.12	The sympathetic and parasympathetic systems play complementary roles in regulating body physiology	351	9.4	Expression of <i>Fru<sup>M</sup></i> in females is sufficient to produce most aspects of male courtship behavior	380	
8.13	The autonomic nervous system is a multilayered regulatory system	353	9.5	Activity of <i>Fru<sup>M</sup></i> neurons promotes male courtship behavior	381	
8.14	The hypothalamus regulates diverse basic body functions via homeostasis and hormone secretion	354	9.6	<i>Fru<sup>M</sup></i> sensory neurons process mating-related sensory cues	382	
<b>HOW IS EATING REGULATED?</b>			356	9.7	<i>Fru<sup>M</sup></i> central neurons integrate sensory information and coordinate the behavioral sequence	384
8.15	Hypothalamic lesion and parabiosis experiments suggested that eating is inhibited by a negative feedback signal from the body	356	9.8	<i>Fru<sup>M</sup></i> neurons in the ventral nerve cord regulate mating-related behavioral output	385	
8.16	Studies of mouse mutants led to the discovery of the leptin feedback signal from adipose tissues	357	9.9	<i>Fru<sup>M</sup></i> -equivalent neurons in females promote female receptivity to courtship	386	
8.17	POMC and AgRP neurons in the arcuate nucleus are central regulators of eating	358	9.10	<i>Fru<sup>M</sup></i> and <i>Doublesex</i> ( <i>Dsx</i> ) regulate sexually dimorphic programmed cell death	386	
8.18	Multiple feedback signals and neural pathways act in concert to regulate eating	360	9.11	<i>Dsx</i> and <i>Fru<sup>M</sup></i> control sexually dimorphic neuronal wiring	389	
<b>HOW ARE CIRCADIAN RHYTHMS AND SLEEP REGULATED?</b>			362	9.12	Even innate behavior can be modified by experience	390
8.19	Circadian rhythms are driven by an auto-inhibitory transcriptional feedback loop that is conserved from flies to mammals	362	<b>HOW ARE MAMMALIAN SEXUAL BEHAVIORS REGULATED?</b>			390
8.20	Entrainment in flies is accomplished by light-induced degradation of circadian rhythm regulators	365	9.13	The <i>Sry</i> gene on the Y chromosome determines male differentiation via testosterone production	393	
8.21	Pacemaker neurons in the mammalian suprachiasmatic nucleus integrate input and coordinate output	366	9.14	Testosterone and estradiol are the major sex hormones	393	
8.22	Sleep is widespread in the animal kingdom and exhibits characteristic electroencephalogram patterns in mammals	367	9.15	Early exposure to testosterone causes females to exhibit male-typical sexual behavior	395	
8.23	The mammalian sleep-wake cycle is regulated by multiple neurotransmitter and neuropeptide systems	369	9.16	Testosterone exerts its organizational effect mainly through the estrogen receptors in rodents	396	
8.24	Why do we sleep?	372	9.17	Dialogues between the brain and gonads initiate sexual maturation at puberty and maintain sexual activity in adults	396	
<b>SUMMARY</b>			374	9.18	Sex hormones specify sexually dimorphic neuronal numbers by regulating programmed cell death	398
<b>FURTHER READING</b>			375	9.19	Sex hormones also regulate sexually dimorphic neuronal connections	399
<b>Chapter 9</b>				9.20	Sexually dimorphic nuclei define neural pathways from olfactory systems to the hypothalamus	400
<b>Sexual Behavior</b>			<b>377</b>	9.21	Whereas the main olfactory system is essential for mating, the accessory olfactory system discriminates sex partners in mice	401
<b>HOW DO GENES SPECIFY SEXUAL BEHAVIOR IN THE FLY?</b>			378	9.22	The same neuronal population can control multiple behaviors in females and males	402
9.1	<i>Drosophila</i> courtship follows a stereotyped ritual that is instinctive	378				

9.23	Parental behavior is activated by mating and regulated by specific populations of hypothalamic neurons	405	WHAT IS THE RELATIONSHIP BETWEEN LEARNING AND SYNAPTIC PLASTICITY?	434	
9.24	Two neuropeptides, oxytocin and vasopressin, regulate pair bonding and parental behavior	407	10.14	Animals exhibit many forms of learning	434
SUMMARY		410	10.15	Habituation and sensitization in <i>Aplysia</i> are mediated by changes of synaptic strength	437
FURTHER READING		412	10.16	Both short-term and long-term memory in <i>Aplysia</i> engage cAMP signaling	439
<b>Chapter 10</b>			10.17	Olfactory conditioning in <i>Drosophila</i> requires cAMP signaling	441
<b>Memory, Learning, and Synaptic Plasticity</b>		<b>415</b>	10.18	<i>Drosophila</i> mushroom body neurons are the site of CS-US convergence for olfactory conditioning	442
PRELUDE: WHAT IS MEMORY, AND HOW IS IT ACQUIRED BY LEARNING?		415	10.19	In rodents, spatial learning and memory depend on the hippocampus	446
10.1	Memory can be explicit or implicit, short-term, or long-term: Insights from amnesic patients	415	10.20	Many manipulations that alter hippocampal LTP also alter spatial memory	447
10.2	Hypothesis I: Memory is stored as strengths of synaptic connections in neural circuits	417	10.21	From correlation to causation: the synaptic weight matrix hypothesis revisited	449
10.3	Hypothesis II: Learning modifies the strengths of synaptic connections	420	WHERE DOES LEARNING OCCUR, AND WHERE IS MEMORY STORED IN THE BRAIN?	451	
HOW IS SYNAPTIC PLASTICITY ACHIEVED?		420	10.22	The neocortex contributes to long-term storage of explicit memory	451
10.4	Long-term potentiation (LTP) of synaptic efficacy can be induced by high-frequency stimulation	421	10.23	The amygdala plays a central role in fear conditioning	454
10.5	LTP at the hippocampal CA3 → CA1 synapse exhibits input specificity, cooperativity, and associativity	421	10.24	Dopamine plays a key role in reward-based learning	456
10.6	The NMDA receptor is a coincidence detector for LTP induction	423	10.25	Early experience can leave behind long-lasting memory traces to facilitate adult learning	459
10.7	Recruitment of AMPA receptors to the postsynaptic surface is the predominant mechanism of LTP expression	423	SUMMARY	463	
10.8	CaMKII auto-phosphorylation creates a molecular memory that links LTP induction and expression	425	FURTHER READING	464	
10.9	Long-term depression weakens synaptic efficacy	426	<b>Chapter 11</b>		
10.10	Spike-timing-dependent plasticity can adjust synaptic efficacy bidirectionally	428	<b>Brain Disorders</b>	<b>467</b>	
10.11	Dendritic integration in the postsynaptic neuron also contributes to synaptic plasticity	428	ALZHEIMER'S DISEASE AND OTHER NEURODEGENERATIVE DISEASES	467	
10.12	Postsynaptic cells can produce retrograde messengers to regulate neurotransmitter release by their presynaptic partners	429	11.1	Alzheimer's disease is defined by brain deposition of numerous amyloid plaques and neurofibrillary tangles	468
10.13	Long-lasting changes of connection strengths involve formation of new synapses	431	11.2	Amyloid plaques mainly consist of aggregates of proteolytic fragments of the amyloid precursor protein (APP)	469
			11.3	Mutations in human APP and $\gamma$ -secretase cause early-onset familial Alzheimer's disease	470
			11.4	Animal models offer crucial tools to investigate pathogenic mechanisms	472
			11.5	An apolipoprotein E (ApoE) variant is a major risk factor for Alzheimer's disease	473

11.6	Microglia dysfunction contributes to late-onset Alzheimer's disease	474	11.26	Synaptic dysfunction is a common cellular mechanism that underlies neurodevelopmental and psychiatric disorders	506
11.7	How can we treat Alzheimer's disease?	475	11.27	Studies of brain disorders and basic neurobiology research advance each other	507
11.8	Prion diseases are caused by propagation of protein-induced protein conformational change	477		<b>SUMMARY</b>	510
11.9	Aggregation of misfolded proteins is associated with many neurodegenerative diseases	479		<b>FURTHER READING</b>	511
11.10	Parkinson's disease results from death of substantia nigra dopamine neurons	480		<b>Chapter 12</b>	
11.11	$\alpha$ -Synuclein aggregation and spread are prominent features of Parkinson's pathology	480		<b>Evolution of the Nervous System</b>	<b>513</b>
11.12	Mitochondrial dysfunction is central to the pathogenesis of Parkinson's disease	482		<b>GENERAL CONCEPTS AND APPROACHES IN EVOLUTIONARY ANALYSIS</b>	514
11.13	Treating Parkinson's disease: L-dopa, deep brain stimulation, and cell-replacement therapy	483	12.1	Phylogenetic trees relate all living organisms in a historical context	515
11.14	The various neurodegenerative diseases have common themes and exhibit unique properties	487	12.2	Cladistic analysis distinguishes processes of evolutionary change	517
	<b>PSYCHIATRIC DISORDERS</b>	487	12.3	Gene duplication, diversification, loss, and shuffling provide rich substrates for natural selection	519
11.15	Schizophrenia can be partially alleviated by drugs that interfere with dopamine function	488	12.4	Altering patterns of gene expression is an important mechanism for evolutionary change	520
11.16	Mood disorders have been treated by manipulating monoamine neurotransmitter metabolism	490	12.5	Natural selection can act on multiple levels in the developing and adult nervous systems to enhance fitness	521
11.17	Modulating GABAergic inhibition can alleviate symptoms of anxiety disorders	491		<b>EVOLUTION OF NEURONAL COMMUNICATION</b>	522
11.18	Addictive drugs hijack the brain's reward system by enhancing the action of VTA dopamine neurons	493	12.6	Ion channels appeared sequentially to mediate electrical signaling	523
11.19	Human genetic studies suggest that many genes contribute to psychiatric disorders	495	12.7	Myelination evolved independently in vertebrates and large invertebrates	524
	<b>NEURODEVELOPMENTAL DISORDERS</b>	498	12.8	Synapses likely originated from cell junctions in early metazoans	525
11.20	Intellectual disabilities and autism spectrum disorders are caused by mutations in many genes	499	12.9	Neurotransmitter release mechanisms were co-opted from the secretory process	526
11.21	Rett syndrome is caused by defects in MeCP2, a regulator of global gene expression	500		<b>EVOLUTION OF SENSORY SYSTEMS</b>	527
11.22	MeCP2 acts predominantly in post-mitotic neurons to regulate their maturation and function	502	12.10	G-protein-coupled receptors (GPCRs) are ancient chemosensory receptors in eukaryotes	530
11.23	Restoring MeCP2 expression in adulthood reverses symptoms in a mouse model of Rett syndrome	503	12.11	Chemosensory receptors in animals are predominantly GPCRs	532
11.24	Fragile-X syndrome is caused by loss of an RNA-binding protein that regulates translation	504	12.12	Two distinct families of ligand-gated ion channels cooperate to sense odors in insects	532
11.25	Reducing mGluR signaling ameliorates fragile-X symptoms in animal models	505	12.13	Retinal- and opsin-based light-sensing apparatus evolved independently at least twice	534

12.14	Photoreceptor neurons evolved in two parallel paths	535	13.7	Reverse genetics disrupts pre-designated genes to assess their functions	563
12.15	Diversification of cell types is a crucial step in the evolution of the retinal circuit	538	13.8	RNA interference (RNAi)-mediated knockdown can also be used to assess gene function	567
12.16	Trichromatic color vision in primates originated from variations and duplications of a cone opsin gene	540	13.9	Genetic mosaic analysis can pinpoint which cell is critical for mediating gene action	568
12.17	Introducing an extra cone opsin in dichromatic animals enables superior spectral discrimination	542	13.10	Transgene expression can be controlled in both space and time in transgenic animals	569
<b>EVOLUTION OF NERVOUS SYSTEM STRUCTURE AND DEVELOPMENT</b>		543	13.11	Transgene expression can also be achieved by viral transduction and other transient methods	571
12.18	All bilaterians share a common body plan specified by conserved developmental regulators	544	13.12	Accessing specific neuronal types facilitates functional circuit dissection	572
12.19	Eye development is controlled by evolutionarily conserved transcription factors	546	13.13	Gene expression patterns can be determined by multiple powerful techniques	572
12.20	The mammalian neocortex underwent rapid expansion recently	547	13.14	Genome sequencing reveals connections across species and identifies genetic variations that contribute to diseases	574
12.21	The size of the neocortex can be altered by modifying the mechanisms of neurogenesis	548	<b>ANATOMICAL TECHNIQUES</b>		575
12.22	Cortical area specialization can be shaped by input patterns	550	13.15	Histological analyses reveal the gross organization of the nervous system	575
<b>SUMMARY</b>		553	13.16	Visualizing individual neurons opens new vistas in understanding the nervous system	578
<b>FURTHER READING</b>		555	13.17	Fine structure studies can identify key facets of molecular organization within neurons	579
<b>Chapter 13</b>			13.18	Mapping neuronal projections allows the tracking of information flow across different brain regions	582
<b>Ways of Exploring</b>		<b>557</b>	13.19	Mapping synaptic connections reveals neural circuitry	584
<b>ANIMAL MODELS IN NEUROBIOLOGY RESEARCH</b>		557	<b>RECORDING AND MANIPULATING NEURONAL ACTIVITY</b>		586
13.1	Some invertebrates provide large, identifiable neurons for electrophysiological investigations	557	13.20	Extracellular recordings can detect the firing of individual neurons	587
13.2	<i>Drosophila</i> and <i>C. elegans</i> allow sophisticated genetic manipulations	558	13.21	Intracellular and whole-cell patch recordings can measure synaptic input in addition to firing patterns	589
13.3	Diverse vertebrate animals offer technical ease or special faculties	559	13.22	Optical imaging can measure the activity of many neurons simultaneously	591
13.4	Mice, rats, and nonhuman primates are important models for mammalian neurobiology research	560	13.23	Neuronal inactivation can be used to reveal which neurons are essential for circuit function and behavior	596
13.5	Human studies are facilitated by a long history of medicine and experimental psychology and by the recent genomic revolution	560	13.24	Neuronal activation can establish sufficiency of neuronal activity in circuit function and behavior	598
<b>GENETIC AND MOLECULAR TECHNIQUES</b>		561	13.25	Optogenetics allows control of the activity of genetically targeted neurons with millisecond precision	599
13.6	Forward genetic screens use random mutagenesis to identify genes that control complex biological processes	562			

13.26 Synaptic connections can be mapped by physiological and optogenetic methods	601
BEHAVIORAL ANALYSES	602
13.27 Studying animal behavior in natural environments can reveal behavioral repertoires and their adaptive value	603
13.28 Studying behaviors in highly controlled conditions facilitates investigation of their neural basis	604
13.29 Behavioral assays can be used to evaluate the functions of genes and neurons and to model human brain disorders	606
SUMMARY AND PERSPECTIVES	608
FURTHER READING	610
GLOSSARY	612
INDEX	